



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Insensitivity of tree-ring growth to temperature and precipitation sharpens the puzzle of enhanced pre-eruption NDVI on Mt. Etna (Italy)

Seiler, Ruedi ; Kirchner, James W ; Krusic, Paul J ; Tognetti, Roberto ; Houlié, Nicolas ; Andronico, Daniele ; Cullotta, Sebastiano ; Egli, Markus ; D'Arrigo, Rosanne ; Cherubini, Paolo

Abstract: On Mt. Etna (Italy), an enhanced Normalized Difference in Vegetation Index (NDVI) signature was detected in the summers of 2001 and 2002 along a distinct line where, in November 2002, a flank eruption subsequently occurred. These observations suggest that pre-eruptive volcanic activity may have enhanced photosynthesis along the future eruptive fissure. If a direct relation between NDVI and future volcanic eruptions could be established, it would provide a straightforward and low-cost method for early detection of upcoming eruptions. However, it is unclear if, or to what extent, the observed enhancement of NDVI can be attributed to volcanic activity prior to the subsequent eruption. We consequently aimed at determining whether an increase in ambient temperature or additional water availability owing to the rise of magma and degassing of water vapour prior to the eruption could have increased photosynthesis of Mt. Etna's trees. Using dendro-climatic analyses we quantified the sensitivity of tree ring widths to temperature and precipitation at high elevation stands on Mt. Etna. Our findings suggest that tree growth at high elevation on Mt. Etna is weakly influenced by climate, and that neither an increase in water availability nor an increase in temperature induced by pre-eruptive activity is a plausible mechanism for enhanced photosynthesis before the 2002/2003 flank eruption. Our findings thus imply that other, yet unknown, factors must be sought as causes of the pre-eruption enhancement of NDVI on Mt. Etna.

DOI: <https://doi.org/10.1371/journal.pone.0169297>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-146034>

Journal Article

Accepted Version

Originally published at:

Seiler, Ruedi; Kirchner, James W; Krusic, Paul J; Tognetti, Roberto; Houlié, Nicolas; Andronico, Daniele; Cullotta, Sebastiano; Egli, Markus; D'Arrigo, Rosanne; Cherubini, Paolo (2017). Insensitivity of tree-ring growth to temperature and precipitation sharpens the puzzle of enhanced pre-eruption NDVI on Mt. Etna (Italy). *PLoS ONE*, 12(1):0169297.

DOI: <https://doi.org/10.1371/journal.pone.0169297>

Insensitivity of tree-ring growth to temperature and precipitation sharpens the puzzle of enhanced pre-eruption NDVI on Mt. Etna (Italy)

short title: Climatic response of trees on Mt. Etna

Ruedi Seiler^{1,2}, James W. Kirchner^{1,3}, Paul J. Krusic^{4,5}, Roberto Tognetti⁶, Nicolas Houlié⁷, Daniele Andronico⁸, Sebastiano Cullotta⁹, Markus Egli², Rosanne D'Arrigo¹⁰ and Paolo Cherubini^{1,2,10}

¹Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

²Department of Geography, University of Zurich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

³Department of Environmental Systems Science, ETH Zurich, CH-8093 Zürich, Switzerland

⁴Navarino Environmental Observatory, Messinia, Greece

⁵Institutionen för Naturgeografi, Stockholm University, Sweden

⁶Dipartimento di Bioscienze e territorio, Università del Molise, Contrada Fonte Lappone, 86090 Pesche, Italy

⁷Department of Earth Sciences, ETH Zurich, CH-8093 Zürich, Switzerland

⁸Osservatorio Etneo, INGV, Sezione di Catania, Italy

⁹Università di Palermo, Palermo, Italy

¹⁰Lamont-Doherty Earth Observatory, Palisades, New York, USA

Abstract

On Mt. Etna (Italy), an enhanced Normalized Difference in Vegetation Index (NDVI) signature was detected in the summers of 2001 and 2002 along a distinct line where, in November 2002, a flank eruption subsequently occurred. These observations suggest that pre-eruptive volcanic activity may

28 have enhanced photosynthesis along the future eruptive fissure. If a direct relation between NDVI
29 and future volcanic eruptions could be established, it would provide a straightforward and low-cost
30 method for early detection of upcoming eruptions. However, it is unclear if, or to what extent, the
31 observed enhancement of NDVI can be attributed to volcanic activity prior to the subsequent
32 eruption. We consequently aimed at determining whether an increase in ambient temperature or
33 additional water availability owing to the rise of magma and degassing of water vapour prior to the
34 eruption could have increased photosynthesis of Mt. Etna's trees. Using dendro-climatic analyses
35 we quantified the sensitivity of tree ring widths to temperature and precipitation at high elevation
36 stands on Mt. Etna. Our findings suggest that tree growth at high elevation on Mt. Etna is weakly
37 influenced by climate, and that neither an increase in water availability nor an increase in
38 temperature induced by pre-eruptive activity is a plausible mechanism for enhanced photosynthesis
39 before the 2002/2003 flank eruption. Our findings thus imply that other, yet unknown, factors must
40 be sought as causes of the pre-eruption enhancement of NDVI on Mt. Etna.

41

42

43 **Introduction**

44 Early detection of precursors to volcanic eruptions is important in preventing major damage and
45 loss of life. To date, these precursors have mainly included seismic, geochemical, petrographic,
46 ground deformation and gravimetric changes that are used to assess volcanic activity shortly before
47 eruptions (e.g., McNutt, 1996; Battaglia et al., 1999; Williams-Jones & Rymer, 2002; Sherburn et
48 al., 2007; Hooper et al., 2012; Sicali et al., 2015). Surface deformation, small earthquakes, and
49 release of volcanic gases are typically triggered by the ascent of magma in volcanoes (Sparks et al.,
50 2012). Volcanic monitoring by remote sensing includes the acquisition of different geochemical and
51 geophysical parameters which record volcanic processes, such as gas emissions and hydrological

52 variations, over time (e.g., Jong, 1998; McNutt et al., 2000; Andronico et al., 2009). Using remote
53 sensing data, Houlié et al. (2006) observed an increased Normalized Difference Vegetation Index
54 (NDVI) along subsequent eruptive fissures on Mt. Nyiragongo (Congo) and Mt. Etna (Italy), as
55 early as two years prior to eruptions of both volcanoes. NDVI is closely associated with the amount
56 of photosynthetically active radiation intercepted by vegetation, and thus with both the spatial
57 coverage of green biomass and the chlorophyll content in leaves (Gamon et al., 1995; Goetz et al.,
58 2005). On Mt. Etna, the enhanced NDVI signal (Houlié et al., 2006) was detected along a narrow
59 line on the northeastern flank; this line later developed into an eruptive fissure during the 2002/2003
60 flank eruption, suggesting that enhanced photosynthesis may be related to a coming volcanic
61 eruption.

62 The observed NDVI signal raises the question of whether, and how, eruptive precursor activity
63 could influence photosynthetic rates. A comparison of tree growth with environmental parameters is
64 necessary to estimate their influence on tree growth and to assess the potential contribution from
65 volcanic activity. Increased photosynthesis may be induced by a number of environmental factors
66 that are affected by pre-eruptive volcanic processes, but the most probable are an increase in heat or
67 water availability associated with volcanic degassing (e.g., Andronico et al., 2005; Aiuppa et al.,
68 2007; Shinohara, 2008).

69 Trees in temperate climates form annual growth rings, and variations in their tree-ring
70 characteristics (width and density) reflect changes in the environmental conditions in which they
71 grow. At high elevations and high latitudes, where the limiting factor is summer air temperature,
72 tree growth is typically enhanced during warm summers (e.g., Buntgen et al., 2006; Briffa et al.,
73 2013). Conversely, in semi-arid ecosystems, such as in Mediterranean lowlands, growth is primarily
74 regulated by precipitation and is enhanced during wet years (Cherubini et al., 2003; Gea-Izquierdo

et al., 2011). Consequently, tree rings, often used as indicators of photosynthetic rates (Dobbertin, 2005), also serve as useful proxies for climate (Fritts, 1976; Begum et al., 2013).

The Mediterranean region is characterized by hot and dry summers and mild, humid winters (Köppen, 1923; Walter & Lieth, 1960). Maximum rainfall occurs predominantly in autumn and sometimes during winter (Bolle et al., 2006). Precipitation minima and temperature maxima coincide with the period of most intense solar radiation, limiting water availability during the summer season (e.g., Ma et al., 2007). High rainfall variability over the year greatly affects drought severity and hampers growth (Cotrufo et al., 2011). Rainfall during spring is the most important factor influencing tree growth and vegetation activity of Mediterranean forests, particularly at more xeric, low-elevation sites, as also shown by remote-sensing-based model simulations and tree-ring-based growth analyses (Maselli et al., 2014). In more temperate high-elevation conditions, drought often has a minor impact on tree growth because precipitation is less limiting. In the high-elevation forests on Mt. Etna where Houlié et al. (2006) detected the increase in NDVI prior to eruption, tree growth might be enhanced by increased air temperature during the vegetation period or, given the southern latitude, by increased water availability.

Here we analyse the relationships between ring width indices of *Pinus nigra* J.F. Arnold and monthly precipitation and air temperature, and compare our ring-width series from Mt. Etna with series from trees growing at similar elevations in Calabria, a region located at a similar latitude on the Italian peninsula without the direct influence of volcanic activity. Our hypotheses are that i) ascending magma led to an increase in local ambient air and soil temperature which positively influenced photosynthesis rates and tree growth, and that ii) water vapour from volcanic degassing locally provided additional humidity/moisture/water which became available to trees influencing photosynthesis rates and tree growth. To address these issues we assess to what extent tree-ring growth at the highest elevations on Mt. Etna is influenced by climate, i.e. air temperature and

99 precipitation, to indirectly determine i) whether an increase in temperature caused by an incipient
100 volcanic eruption (e.g., Bonneville & Gouze, 1992; Andronico et al., 2005) would likely induce
101 higher photosynthetic productivity, and ii) whether, at specific locations close to rift zones,
102 additional water availability induced by degassing of water vapour associated with the rise of
103 magma prior to an eruption (water is the most abundant component in volcanic gas, Shinohara,
104 2008) would likely increase the photosynthetic capacity of Mt. Etna's trees (see Houlié et al., 2006).

105

106

107 **Materials and Methods**

108 *Study area*

109 Mt. Etna is a stratovolcano situated in the northeastern part of Sicily. With an area of approximately
110 1600 km² and a summit elevation of roughly 3330 m a.s.l., Mt. Etna is an isolated high mountain
111 exposed to air masses from the Mediterranean Sea. The climate on Mt. Etna is strongly maritime on
112 the eastern flank (Poli Marchese & Grillo, 2004; Branca et al., 2008), with drier conditions on its
113 western flank (Burga & Klötzli, 2004). The slopes are characterized by lava flows of different ages
114 (Doglioni et al., 2002). Most of the lower elevations, being especially fertile, have been settled and
115 used for agriculture for thousands of years. The higher elevations, from 1000-1600 m a.s.l., are
116 dominated by European beech (*Fagus sylvatica* L.), and from 1600 m to treeline (~2000 m a.s.l.) by
117 Corsican black pine (*P. nigra*). Though the treeline climatically determined at such latitudes would
118 otherwise be higher (Hermes, 1955; Körner, 1998), eruption-induced wildfires and the lack of soils
119 on lava flows hinder its uphill development (Certini et al., 2001; Dazzi, 2007; Egli et al., 2012).

120 Besides volcanic eruptions and lava flows on Mt. Etna, other volcanic processes, such as degassing
121 through small vents, are also present but difficult to quantify (Allard, 1997). The soils of Mt. Etna
122 are primarily classified as Regosols, Eutric or Dystric Cambisols and (Mollic) Andosols. The

characteristics of these soils predominantly depend on the surface age of the lava flow and volcanic deposits from which they have developed (Dazzi, 2007; Lulli, 2007; Egli et al., 2008). In general, soils at intermediate to high elevation on Mt. Etna (i.e. above 800 m a.s.l.) are mostly described as Andisols with a sandy loam texture, vitric characteristics, an udic moisture regime (Egli et al., 2012) and good water holding capacity (Maeda et al., 1977). However, less mature, young soils on fresh lava flows may be less developed resulting in lower water holding capacity.

The forests around the flanks of Mt. Etna are greatly affected by both natural and anthropogenic disturbances, such as wildfires, lava flows, avalanches and logging. At the lowest elevations, from the plains up to 900 m a.s.l., agricultural crops and orchards, e.g., orange, lemon, almond, pistachio, and chestnut, are found. Only a few forest stands, mainly at the highest elevations on the northern or northeastern side of the mountain, are undisturbed (Poli Marchese & Grillo, 2004). Meteorological station data from Linguaglossa (530 m a.s.l., 15°08'42" E, 37°50'27" N, timespan: 1893-2004) based on daily temperatures and precipitation measurements give an average annual temperature of 18°C and a total annual precipitation of 1400 mm. Additionally, monthly temperature averages in winter are above zero at all stations.

138

139 *Sampling*

In total, we sampled 143 trees (*P. nigra*), with permission issued by the local forest authorities (Corpo Forestale della Regione Siciliana, Distaccamento di Bronte, Piazza Cadorna 11, I-95034 Bronte, Catania, Italy), at four high-elevation (1500 to 1900 m a.s.l.) forest sites on the northeastern and western slopes of Mt. Etna (Fig 1): 52 trees growing close to the 2002/2003 eruptive fissure (Group 1), 27 trees growing close to the 1928 eruptive fissure (Group 2), 38 trees growing in the same elevation band but far from any obvious fissures (Group 3), and 26 trees growing close to the 1974 eruptive fissure (Group 4). All sites are located at a comparable elevation and slope with NNE

147 aspect except for group 4, which is located on the western flank on a western-aspect slope. Apart
148 from that, there were no evident differences between the four sites in terms of forest-stand density,
149 composition of tree species, topography and slope (about 14%). From each tree, two 0.5 cm
150 diameter cores were taken orthogonally with respect to each other using a corer with a three-
151 threaded auger by Haglof (Haglof Inc., Sweden), wrapped in paper and transported to the
152 laboratory. All samples were mounted on wooden supports and cut following the technique
153 suggested by Gärtner et al. (2015), using a microtome at an angle of roughly 30° to the radial axis
154 of the tree to prevent core breakage. For later comparisons, three ring-width chronologies, located
155 close to Mt. Etna, derived from coniferous trees uninfluenced by Mt. Etna growing at similarly high
156 elevations in Calabria (Gambarie, Monte Pollino and Sierra da Crispo) were retrieved from the
157 International Tree-Ring Databank (ITRDB, NOAA, U.S.A.)

158

159 **Fig 1. Map of sample locations.**

160 Mt. Etna sample sites (Group 1-4) on the northeastern and western slopes at an elevation range from
161 1600 to 1850 m a.s.l. indicating the location where samples were taken and the location of the
162 meteorological stations.

163

164 *Ring-width measurements*

165 All ring widths were measured to the nearest 0.01 mm using a Leica Wild M32 binocular
166 microscope (Leica, Germany) with 25-50x magnification, coupled to a LINTAB measuring table
167 and computer with TSAPwin (Time Series Analysis Program) software (RinnTech, Heidelberg,
168 Germany). Core measurements were visually crossdated against each other and any inconsistencies,
169 if found, were eliminated. Subsequent crossdating of the single-tree chronologies with their
170 respective mean site chronology by visual and statistical measures was performed using TSAPwin

171 and COFECHA (50-year segments with a 25-year overlap) (Holmes, 1983; Grissino-Mayer et al.,
172 2001). Since the sampling dates of all trees were known, crossdating was primarily used to ensure
173 that prominent tree-ring patterns were not shifted between trees and no rings were missing.

174

175 *Meteorological data*

176 We used monthly precipitation and air temperature data recorded at three meteorological stations on
177 Mt. Etna: Floresta (1275 m a.s.l., 14°54'31" E, 37°59'15" N, Timespan: 1924-2004), Linguaglossa
178 (530 m a.s.l., 15°08'42" E, 37°50'27" N, Timespan: 1893-2004) and Taormina (248 m a.s.l.,
179 15°17'34" E, 37°17'34" N, Timespan: 1906-2004). Although longer records are available for
180 Linguaglossa and Taormina, we only used data from 1924-2004 at all three sites so that they could
181 be compared over a common period. In addition, interpolated monthly temperature, precipitation,
182 cloud cover and Palmer Drought Severity Index (PDSI) data for the Mt. Etna region and Calabria
183 from the Climatic Research Unit, University of East Anglia, Norwich, U.K. (CRU; Mitchell et al.,
184 2004) were compared to the above-mentioned station data and the tree-ring data (Griggs et al.,
185 2007; Carrer et al., 2010). Opposed to temperature and precipitation data, the PDSI incorporates
186 both temperature and precipitation, representing long-term drought taking prior months' condition
187 into account. On the other hand, delayed water runoff from snow during spring is not accounted for
188 in the index (e.g., Lüdeke et al., 1996). Correlations between the data recorded at the meteorological
189 stations and the interpolated datasets were calculated, to assess whether the interpolated data could
190 be used to further analyse relationships between climate and tree growth.

191

192 *Data analysis*

193 All raw measurement series were standardized using the program ARSTAN
194 (<http://www.ldeo.columbia.edu/tree-ring-laboratory>) by applying 30-year spline detrending

195 combined with a variance stabilization, to remove the age trend and produce detrended ring-width
196 indices (Holmes, 1983; Cook, 1985; Büntgen et al., 2005). All chronologies were used individually
197 to analyse the relationships between climate and growth at different sampling sites.

198

199 We performed correlation analysis and response function modelling to quantify the influence of
200 climate on tree growth (Cook et al., 1990). We tested the statistical significance of temperature,
201 precipitation, cloud cover and Palmer Drought Severity Index (PDSI) (e.g., García-Suárez et al.,
202 2009; Cai et al., 2014) in different months and seasonal combinations of monthly values, including
203 prior-year values, using Spearman rank correlation. Linear regression, as described by Tomé &
204 Miranda (2004), was used to remove long-term trends in the meteorological data and avoid
205 artificially inflating correlation values. Based on results from simple Spearman rank correlations
206 between all the Mt. Etna chronologies and monthly meteorological data, we built "Visual
207 Regression Models" (VRM) which were defined as standard multiple linear regression models
208 including statistically significant ($p < 0.05$) monthly variables or monthly groupings.

209 In addition to VRM, Stepwise Linear Regression Modelling (SLRM), based on the Akaike
210 Information Criterion (AIC) and using a forward-backward approach (Cook & Pederson, 2011),
211 was used to identify those climate variables that explained the greatest variance in each chronology
212 on Mt. Etna. The SLRMs were based on monthly variables and groupings that defined the spring
213 and summer seasons. For all models we only used variables that did not overlap in time (e.g.,
214 precipitation in May and precipitation in spring would not be used together because both contain
215 May precipitation values).

216 The models (SLRM) were tested for collinearity among explanatory variables using variance
217 inflation factor (VIF) analysis (Kutner et al., 2005). VIF values were all lower than 4, implying a
218 lack of strong collinearity among explanatory variables (O'Brian, 2007). This led to selecting the

219 final climate models based on the AIC.

220 We compared how well the two model types (VRM and SLRM) explained the climate – ring width
221 relationship:

222 1) Qualitative differences between the two model types included differences in monthly parameters
223 used in the models.

224 2) Quantitative differences, comparing adjusted- R^2 , show the increase/decrease of explanatory
225 power from one model type to another.

226 To test the reliability of the models, as well as the degree of overfitting due to including too many
227 variables during the stepwise model selection process, we divided the timespan covered by both
228 meteorological and ring-width measurements at Mt. Etna (1924-2004) and Calabria (1924-1980)
229 into two segments (Todaro et al., 2007; Griggs et al., 2014). The models were run on both segments
230 to compare differences over time. In addition, the two segments were used for both forward- and
231 backward validation to quantify model robustness over time (Cook & Kariukstis, 1990).

232 On Mt. Etna, forward validation used a model based on the time segment 1925-1964 to predict tree
233 growth during 1965-2004, and backward validation used a model based on the latter time segment
234 to reconstruct 1925-1964 tree growth. For the Calabria chronologies we used the same validation
235 procedure based on the two time segments from 1925-1951 and 1952-1980. AIC was used to test
236 for overfitting.

237 We calculated how the explanatory power (R^2) varied through time by applying our SLRMs to a 15-
238 year moving window with 14 years overlap to identify periods of higher and lower correlations
239 between ring-width and our climate models (Carrer et al., 2010).

240

241

242 **Results**

243 The Mt. Etna chronologies show higher variability in their ring-width patterns than the chronologies
 244 from Calabria (Fig 2). The four raw ring-width chronologies (Group1-4) from Mt. Etna display
 245 growth patterns that are strongly influenced by the establishment of new generations of trees.
 246 Young trees, most germinating after 1950, greatly increase the mean ring-width, producing a clear
 247 age trend (Fig 2). Tree age ranges from 55 to 122 years on Mt. Etna, and from 128 to 299 years in
 248 Calabria. Descriptive chronology statistics are given in Table 1.

249

250 Fig 2. **Chronologies and sample replication.**

251 Average chronologies (black lines) and sample replication of all samples (grey area) of Mt. Etna
 252 (Group 1-4) and Calabria (Gambarie, Monte Pollino and Sierra da Crispo).

253

| | no. of series | total length | ser.interc. | mean length | elevation | species | |
|----------|------------------|--------------|-------------|-------------|-----------|---------|--------------------------|
| Mt. Etna | Group 1 | 104 | 229 | 0.498 | 97 | 1850 | <i>Pinus nigra</i> |
| | Group 2 | 54 | 121 | 0.577 | 67.1 | 1700 | <i>Pinus nigra</i> |
| | Group 3 | 76 | 195 | 0.529 | 82.7 | 1600 | <i>Pinus nigra</i> |
| | Group 4 | 52 | 130 | 0.597 | 95 | 1670 | <i>Pinus nigra</i> |
| Calabria | Gambarie | 26 | 191 | 0.344 | 130.7 | 1850 | <i>Abies alba</i> |
| | Monte Pollino | 24 | 181 | 0.403 | 128.2 | 1720 | <i>Abies alba</i> |
| | Sierra da Crispo | 22 | 540 | 0.421 | 299.1 | 2000 | <i>Pinus leucodermis</i> |

254

255 Table 1. **Sample overview information.**

256 Descriptive statistics of sample chronologies (Group 1-4) from Mt. Etna, and the chronologies from
 257 Calabria (Gambarie, Monte Pollino and Sierra da Crispo) displaying number of series (core-series),

total length (years) of group chronologies, series intercorrelation (measure of common growth signal in the chronology), mean sample length (years), elevation (m a.s.l.) and species.

The monthly meteorological data show stronger inter-station correlations for air temperature (Spearman correlation coefficients up to $r = 0.86$; $p < 0.01$) than for precipitation (up to $r = 0.83$; $p < 0.01$), as usually reported in the literature (e.g., Graumlich, 1987; Griggs et al., 2007). In addition, we find that instrumental station data also correlates significantly with the interpolated CRU data (highest Spearman $r = 0.67$ to 0.87 for temperature; $r = 0.67$ to 0.78 for precipitation; $p < 0.01$). Given these correlations, comparable to those found in previous studies (Griggs et al., 2007), we used the interpolated data to assess the climate influence on tree growth.

In general, total summer precipitation produces similarly high correlation values with tree growth as single monthly variables in the same season. Average spring and summer temperature and total spring precipitation produce generally lower correlations than single months during the same season (Table 2). Prior-year precipitation and temperature were also considered but results are not reported because the current year's correlations are higher. Cloud coverage is not significantly correlated with tree growth. In contrast to all other sites, tree growth at Group 3 exhibits a significant negative correlation with PDSI from April to December (results not shown).

| | Mt. Etna | | | | Calabria | | |
|---|------------------|-----------------|------------------|---------|------------------|-----------------|------------------|
| SPEARMAN rank correlations | Group 1 | Group 2 | Group 3 | Group 4 | Gambarie | Monte Pollino | Sierra da Crispo |
| T February | 0.176 | 0.189 | 0.217 | 0.055 | 0.209 | 0.184 | 0.114 |
| T March | ** 0.427 | * 0.265 | ** 0.327 | 0.197 | 0.152 | 0.163 | 0.018 |
| T April | 0.144 | -0.109 | 0.064 | 0.169 | 0.148 | ** 0.490 | 0.203 |
| T May | -0.106 | -0.213 | -0.15 | -0.033 | -0.018 | * 0.271 | 0.061 |
| T avg. spring | ** 0.337 | 0.197 | * 0.284 | 0.14 | 0.21 | ** 0.348 | 0.11 |
| T June | 0.035 | -0.094 | -0.004 | 0.172 | * -0.283 | -0.019 | -0.068 |
| T July | -0.158 | * -0.223 | -0.136 | 0.06 | * -0.296 | -0.068 | 0.026 |
| T August | ** -0.326 | -0.182 | ** -0.397 | -0.032 | ** -0.408 | -0.05 | -0.131 |

| | | | | | | | |
|------------------|-----------------|-----------------|------------------|----------------|------------------|-----------------|--------|
| T September | -0.117 | -0.147 | -0.198 | -0.01 | -0.178 | 0.026 | 0.117 |
| T avg. summer | -0.172 | -0.206 | -0.203 | 0.088 | ** -0.375 | -0.07 | -0.084 |
| prior P December | -0.16 | * -0.261 | ** -0.304 | -0.028 | 0.087 | 0.181 | 0.111 |
| P January | 0.131 | -0.095 | 0.091 | 0.021 | -0.137 | -0.018 | 0.059 |
| prior P winter | 0.027 | * -0.251 | -0.078 | 0.043 | 0.03 | 0.184 | 0.182 |
| P February | -0.048 | -0.113 | -0.159 | -0.125 | -0.057 | -0.128 | 0.029 |
| P March | * -0.279 | -0.151 | -0.12 | -0.079 | -0.007 | -0.001 | 0.044 |
| P April | 0.009 | 0.121 | 0.15 | 0.103 | -0.22 | * -0.279 | -0.031 |
| P May | 0.185 | 0.088 | * 0.227 | 0.025 | 0.209 | -0.026 | -0.179 |
| P tot. spring | * -0.228 | -0.104 | -0.107 | -0.095 | -0.19 | * -0.308 | -0.015 |
| P June | 0.114 | 0.03 | * 0.236 | -0.117 | * 0.279 | 0.138 | 0.076 |
| P July | * 0.279 | 0.092 | * 0.222 | * 0.225 | * 0.303 | ** 0.416 | 0.186 |
| P August | 0.165 | -0.043 | 0.209 | 0.09 | 0.083 | 0.083 | 0.136 |
| P tot. summer | * 0.271 | 0.09 | ** 0.346 | 0.118 | * 0.325 | * 0.314 | 0.24 |

276

277 Table 2. **Climate-ring width correlation statistics.**

278 Spearman rank correlations between climate variables and detrended ring width from Mt. Etna
279 chronologies (Group 1-4) and from Calabria chronologies (Gambarie, Monte Pollino and Sierra da
280 Crispo), where P = precipitation, T = temperature, tot. = total amount of precipitation, avg. =
281 average temperature, and prior = prior year. Values printed in bold are statistically significant with
282 (* = $p < 0.05$, ** = $p < 0.01$, two-tailed). The significance threshold at Mt. Etna is lower than in
283 Calabria ($r = 0.222$ vs. $r = 0.271$, respectively) because the climate and tree ring records overlap for
284 longer on Mt. Etna than in Calabria (81 vs. 57 years, respectively).

285

286 In Calabria the Gambarie chronology is primarily sensitive to summer temperature and summer
287 precipitation (Table 2). The Monte Pollino chronology responds more to spring temperatures as
288 well as spring and summer precipitation, and the chronology from Sierra da Crispo is not
289 significantly correlated with any climate variables. The Mt. Etna chronologies correlate with spring
290 and summer temperatures (Groups 1-3), as well as spring and summer precipitation (Groups 1 and
291 3). Group 4 has no significant correlations with climate. Over all months and seasons the Mt. Etna

292 (Groups 1-4) chronologies are, on average, less strongly correlated with temperature and
293 precipitation than the Calabria chronologies are. The difference in the (absolute value) strength of
294 the correlations between Mt. Etna and Calabria was not, however, statistically significant (Student
295 *t*-test).

296 Overall we observe similar responses in the chronologies from Mt. Etna and in those from Calabria
297 (Table 2 and Fig 3). The raw numbers of significant correlations between temperature or
298 precipitation and ring width are broadly similar (20 out of 88, or 23% at Mt. Etna, and 14 out of 66,
299 or 21%, in Calabria).

300

301 **Fig 3. Climate - ring-width correlations.**

302 Spearman rank correlation data points of single months and seasonal groupings (red dots) of
303 chronologies from Mt. Etna (top panel) and Calabria (bottom panel) showing the correlation range
304 of each monthly- or seasonal variable with the different group-chronologies; where T =
305 temperature, P = precipitation and p = prior year. Boxplots show the median and lower and upper
306 quartiles, and the whiskers display the minimum and maximum values.

307

308 We used the sign test to evaluate whether the correlations between climate and ring width were
309 positive or negative more often than expected by chance. If we pool both the Mt. Etna and Calabria
310 correlations (Table 2), we see that ring widths are positively correlated with spring temperature (22
311 of 28 month and site combinations; $p < 0.01$ by the two-tailed sign test), negatively correlated with
312 summer temperature (22 of 28 correlations; $p < 0.01$ by the two-tailed sign test), and positively
313 correlated with summer precipitation (18 of 21 correlations; $p < 0.05$ by the two-tailed sign test),
314 but not significantly correlated with spring precipitation (17 negative correlations out of 28 month
315 and site combinations; $p > 0.05$ by the two-tailed sign test). Thus tree growth in these mountain

environments tends to be favoured by warm springs (suggesting that water is not limiting in the springtime) and cool and relatively wet summers.

Comparing the VRM and SLRM modelling results (Table 3), we distinguished between models explaining ring width on Mt. Etna and models explaining ring width in Calabria. SLRM's yielded average R^2 values of 20% (8% to 33%) on Mt. Etna and 26% (13% to 39%) in Calabria, demonstrating that precipitation and temperature are not strongly correlated with tree growth compared to other climatic regions. The regression models include both precipitation and temperature variables from spring and summer. There are, however slight differences between the models (Table 4).

| | Visual Regression Models | | | Stepwise Linear Regression Models | | | Model improvement (%) | |
|-------------------------|--------------------------|-------------|-----------------|-----------------------------------|-------------|----------------|-----------------------|------------|
| | R^2 | adj. R^2 | <i>p-value</i> | R^2 | adj. R^2 | <i>p-value</i> | R^2 | adj. R^2 |
| Group 1 | 0.23 | 0.19 | <0.01 | 0.29 | 0.24 | <0.01 | 6 | 5 |
| Group 2 | 0.09 | 0.04 | 0.13 | 0.09 | 0.06 | 0.03 | n/a | n/a |
| Group 3 | 0.27 | 0.22 | <0.01 | 0.33 | 0.27 | <0.01 | 6 | 5 |
| Group 4 | 0 | -0.01 | 0.96 | 0.08 | 0.06 | 0.03 | n/a | n/a |
| average Mt. Etna | 0.15 | 0.11 | 0.27 | 0.2 | 0.16 | 0.02 | 6 | 5 |
| | | | | | | | | |
| Gambarie | 0.2 | 0.17 | <0.01 | 0.26 | 0.23 | <0.01 | 6 | 6 |
| Monte Pollino | 0.3 | 0.24 | <0.01 | 0.39 | 0.34 | <0.01 | 9 | 10 |
| Sierra da Crispo | - | - | - | 0.13 | 0.1 | 0.02 | n/a | n/a |
| average Calabria | 0.25 | 0.2 | <0.01 | 0.26 | 0.22 | 0.01 | 7.5 | 8 |

326

327 Table 3. **Statistics of climate models.**

328 Overview of model R^2 and adjusted R^2 statistics of the Mt. Etna and Calabria chronology models.

329 Visual Regression Models (VRM) are shown in the left panel, Stepwise Linear Regression Models

(SLRM) in the middle panel and the percentage of "model-improvement" from VRM to SLRM is shown in the right panel.

| | | temperature | | | | | | | | | | precipitation | | | | | | | | | | |
|----------|---------|-------------|-------|-------|-------------|-----|------|------|--------|-------------|----------|---------------|-------------|----------|-------|-------|-------------|-----|------|------|--------|-------------|
| | | February | March | April | avg. spring | May | June | July | August | avg. summer | December | January | tot. winter | February | March | April | tot. spring | May | June | July | August | tot. summer |
| Mt. Etna | Group1 | VRM | X | | | | | X | | | | | | X | | | | | | X | | |
| | | SLRM | X | | | | | | X | | | X | | | | | X | | | | | X |
| | Group 2 | VRM | X | | | | | X | | | X | | e | | | | | | | | | |
| | | SLRM | | | | | X | | | | X | | | | | | | | | | | |
| | Group 3 | VRM | X | | | | | X | | X | | | | | | | | X | | | | X |
| | | SLRM | | | | X | X | | X | e | X | | | | | | | | | | | X |
| | Group 4 | VRM | | | | | | | | | | | | | | | | | | X | | |
| | | SLRM | | | | | | X | | | | | | X | | | | | | | | |

| Calabria | Gambarie | VRM | | | | | | | X | | | | | | | | | | | | | X |
|----------|------------------|------|--|--|---|---|---|--|---|---|---|--|--|--|--|---|--|--|--|---|--|---|
| | | SLRM | | | X | | | | | X | | | | | | | | | | | | |
| | Monte Pollino | VRM | | | X | | X | | | | | | | | | X | | | | X | | |
| | | SLRM | | | X | e | | | | | X | | | | | X | | | | | | X |
| | Sierra da Crispo | VRM | | | | | | | | | | | | | | | | | | | | |
| | | SLRM | | | X | | | | | | | | | | | | | | | | | X |

333

334 Table 4. **Model variables used by climate models.**

335 Climate variables (single months and seasonal groupings) used in VRM and SLRM models.

336 Monthly variables included in the models are designated as X. Due to time overlaps between single

337 months and seasons, variables that were excluded from the models are designated as e.

338

339 We calculated an average improvement in adjusted R^2 from VRM to SLRM of 5% on Mt. Etna and

340 8% in Calabria (Table 3; Fig 4).

341 When comparing VRM and SLRM which were statistically significant ($p < 0.05$), a significant
342 model-improvement ($p < 0.05$) was only obtained with Mt. Etna's Group 1 models. These results
343 show that on Mt. Etna even complex models such as our SLRM are not able to explain tree-growth
344 variability much better, demonstrating that tree growth is further influenced by parameters other
345 than climate which induce additional noise to our ring-width data.

346

347 **Fig 4. Climate model comparison.**

348 Comparisons between individual VRMs (visual regression models) and SLRMs (stepwise linear
349 regression models) revealed only one case (Group 1) where the VRM and the SLRM were both
350 significant ($p < 0.05$) and where a statistically significant model improvement ($p < 0.05$) was
351 calculated. On Mt. Etna, the Group 1 model significantly improved from adjusted $R^2 = 0.19$ (VRM)
352 to adjusted $R^2 = 0.24$ (SLRM). Details are summarized in Table 3.

353

354 When comparing differences over time by running the SLRMs on both time segments separately,
355 on Mt. Etna only two out of eight model-runs on the two time segments led to significant R^2 values
356 ($p < 0.05$), whereas in Calabria the R^2 in four out of six runs was significant (results not shown).

357

358 Mt. Etna model validations demonstrate that the only forward verification resulting in a significant
359 R^2 value ($p < 0.01$) was obtained with the Group 3 model. Forward validation of the Group 1,
360 Group 2 and Group 4 models resulted in statistically non-significant R^2 values. Further, backward
361 validations show that all models calibrated on the second segment show a decrease to non-
362 significant R^2 values when run on the first segment. Out of eight validation runs (forward and
363 backward) on Mt. Etna only one retained significant R^2 values ($p < 0.01$). Including all validation
364 runs (statistically significant and non-significant), forward validation lost 4% (from an average R^2

365 of 0.18 in the first period to 0.14 in the second period) and backward validation lost 37% (from an
366 average R^2 of 0.48 in the second period to 0.11 in the first period) of explained ring-width variance
367 on Mt. Etna.

368 The same calculations for Calabria showed that forward validation (earlier to later time-segment)
369 lost 32% of the explained variance (average R^2 changed from 0.53 to 0.21), while backward
370 validation lost 11% (average R^2 changed from 0.55 to 0.44). We calculated that of all six validation
371 runs (forward and backward) in Calabria, only two backward validations retained significant R^2
372 values ($p < 0.05$).

373 These results show that our models (SLRMs) of tree-ring data on Mt. Etna and in Calabria do not
374 withstand the cross-validation test and demonstrate the importance of validating tree-ring based
375 climate models, especially in regions such as the Mediterranean, where climatic factors are not
376 strongly limiting tree growth.

377 The explanatory power of all models is generally low. The changes in the explanatory power of all
378 models over time are shown in Fig 5. The Mt. Etna models (Groups 1-4) display a wider range of
379 performance with some visual suggestion of a trend of increasing explanatory power over time,
380 while the Calabria models showed less temporal variability.

381

382 Fig 5. **Model strength over time.**

383 Change of SLRM model goodness-of-fit statistics for 15-year moving windows over time, showing
384 a visual suggestion of higher temporal consistency among the Calabria models.

385

386

387 **Discussion**

388 In the Mediterranean region, seasonal drought conditions can persist at low elevations for up to five
389 months and have a strong negative impact on tree growth (Scarascia-Mugnozza et al., 2000;
390 Cherubini et al., 2003), whereas at higher elevations precipitation is usually not a limiting factor
391 (e.g., Maselli et al., 2014). At the treeline in temperate regions, air temperature is generally the
392 major driver of tree growth, being positively correlated with ring width (e.g., Eckstein & Aniol,
393 1981; Schweingruber, 1987; Briffa et al., 1996; Hughes, 2002; Leonelli et al., 2009). Similar
394 correlations have been observed in the Mediterranean mountains as well (Serre-Bachet & Guiot,
395 1987; Büntgen et al., 2012). At lower elevations, higher temperatures reduce tree growth by
396 increasing evaporative demand and drought stress (Campelo et al., 2006; Olivar et al., 2012). Our
397 study showed that tree growth at high elevation on Mt. Etna is not much limited by climatic
398 conditions: the correlations between tree-ring width and meteorological data (monthly precipitation,
399 air temperature, PDSI and cloud coverage) were rather weak, suggesting that an increase in local
400 moisture or temperature caused by pre-eruptive volcanic activity was unlikely to have affected tree
401 growth. On Mt. Etna, the correlations between ring width and meteorological data were weaker
402 than in Calabria, suggesting that the tree-ring/climate relationship on Mt. Etna might be affected by
403 other factors. Stepwise linear regression models (SLRM) explained an average adjusted R^2 of 16%
404 of tree growth on Mt. Etna and 22% in Calabria. By comparison, linear regression models using
405 spring to summer temperature at similar elevations in south-western Anatolia explained up to 51%
406 of ring-width variance (Touchan et al., 2007). Furthermore, climate correlation values on Mt. Etna
407 and in Calabria are lower than those found using response function analyses in the Aegean (Hughes,
408 2001). Tree-growth response to temperature across the Mediterranean region is complex and
409 strongly affected by longitude, elevation (Seim et al., 2012) and the age of the trees studied
410 (Navarro-Cerrillo et al., 2014). The dependence of tree growth on spring temperature may be
411 reduced at sites that, like Mt. Etna and Calabria, are very close to the sea.

412 In general, the ring-width chronologies on Mt. Etna showed higher inter-annual variability than
413 those from Calabria, even though the Mt. Etna chronologies were based on greater numbers of trees
414 with a correspondingly greater averaging of random inter-tree variations. Our analyses suggest that
415 non-climatic factors may give rise to a greater variability in tree-ring growth on Mt. Etna than in
416 Calabria. This argument is supported by the larger differences on Mt. Etna between the regression
417 models fitted to the two time periods.

418 On Mt. Etna, the correlation with summer precipitation ($r = 0.27$ to 0.35) is lower than in previous
419 studies that found that water is the limiting factor in the Mediterranean region (Hughes et al., 2001;
420 Maselli et al., 2004; Touchan et al., 2007; Akkemik et al., 2008; Allard et al., 2008; Griggs et al.,
421 2014). Near treeline on Mt. Etna, which is lower than at other sites at similar latitudes, because not
422 determined by climatic conditions but rather by volcanic activities, other factors, such as higher air
423 humidity with increasing elevation, seem to reduce the influence of summer precipitation on tree
424 growth.

425 The low correlation values between PDSI and ring width on Mt. Etna (maximum $r = 0.29$) confirm
426 that moisture availability is not strongly limiting tree growth. Similar results have been reported by
427 Aloui (1982) for *Pinus halepensis* Mill. in Tunisia and by Tessier (1984) for *Pinus sylvestris* L. in
428 south-eastern France. It is notable that *P. nigra* exhibits a drought-avoidance strategy characterized
429 by efficient stomatal control of transpirational water loss (Lebourgeois et al., 1998). At the same
430 latitudes in Spain, the drought response of *P. nigra* varied along an aridity gradient with the
431 strongest response at the most xeric sites (Camarero et al., 2013). At comparable latitudes and
432 elevations in Calabria, studies on *F. sylvatica* found no correlations between either water use
433 efficiency or basal area increment and an estimated drought index, suggesting a minimal effect of
434 climate on tree growth during the last century (Tognetti et al., 2014), especially when considering
435 the rather udic soil moisture regime at intermediate to high elevations on Mt. Etna (Egli et al.,

2012). No significant differences in soil properties from soils adjacent to eruptive fissures and soils away from such fissures (Reisser, 2014) revealed homogeneity of soils on Mt. Etna.

Due to their high elevation and the subsequent cold, snowy, foggy and humid environmental conditions, the Mt. Etna trees do not appear to be strongly affected by Mediterranean summer drought. The combination of proximity to the sea and high elevation favours persistent seasonal fog and shading by clouds, thus limiting the vapour pressure deficit and raising water use efficiency (Limm et al., 2009). To survive summer droughts, vegetation uses water coming from spring precipitation or, at higher elevation, melting snow, which in some regions can be half of the annual precipitation amount (Renault et al., 2010). At the highest elevations on Mt. Etna, winter precipitation is mainly snowfall and, based on our analyses, is not strongly correlated with tree growth (Table 2). The high porosity of the volcanic soils allows rapid infiltration, making a large fraction of the annual precipitation unavailable to the trees, and making it unlikely that winter or spring precipitation could be stored long enough to significantly alleviate summer drought stress.

Significant positive correlations between ring width and spring temperatures were found, except for Group 4. These results confirm those of previous studies on silver fir in southern Italy (Carrer et al., 2010), and in southwestern Anatolia (Touchan et al., 2014). Based on the positive correlations between March temperatures and ring width on Mt. Etna, high spring temperatures appear to promote an early start of the growing season. Based on the above zero average temperatures during winter measured at all meteorological stations, possible heat discharge from the volcanic fissure in the years before the eruption (e.g., Aubert, 1999) could not have caused such an early start of the growing season. In contrast, pines are able to photosynthesize during winter; thus mild temperatures in late winter may enhance the availability of reserves (non-structural carbohydrates) for allocation to cambial growth in spring. Mild conditions in spring may stimulate cambial dynamics or induce early cambial reactivation, increasing production of early-wood.

460 We found negative correlations between ring width and summer temperature, as previously
461 described by Hughes (2001), Griggs et al. (2007) and Köse et al. (2011; 2013) in Turkey, as well as
462 in north-eastern Greece and the Spanish Pyrenees (Tardif et al., 2003). These correlations may be
463 related to heat waves and the negative effect of drought stress on tree growth. An increase in
464 frequency in hot and dry summers under climate change during the recent past (Christensen et al.,
465 2013) may also increase drought stress indirectly in autumn and further constrain tree growth.

466

467

468 **Conclusions**

469 We conclude that tree growth at the highest elevations on Mt. Etna is not significantly limited by
470 climate. Our samples were taken near Mt Etna's upper treeline, but this treeline is not climatically
471 determined; instead it is defined by volcanic activity and related disturbances such as wildfires.
472 Consequently, climatic influences on tree growth are weaker than would be expected in trees
473 growing at a climatically induced treeline where temperature is the limiting factor (Maselli et al.,
474 2014). At the same time, the Mt. Etna trees are growing at an elevation that is too high to be
475 strongly affected by summer drought as in Mediterranean lowlands (Cherubini et al., 2003). The
476 intermediate elevation between the two extremes (high elevation where temperature is limiting and
477 low elevation where summer drought is limiting) makes it difficult to explain the tree-growth
478 variability using meteorological data. The low sensitivity of tree growth to climate suggests that
479 neither i) an increase of surrounding air temperature caused by heating from magma at shallow
480 depths, nor ii) an increase in water availability induced by pre-eruptive subsurface pressures and
481 water vapour, is likely to have enhanced photosynthesis before the 2002/2003 flank eruption. Thus
482 to explain the NDVI signal previously observed by Houlié et al. (2006), one must search for factors
483 (volcanic or not) other than additional water or heat induced by volcanic activity.

484

485

486 **Acknowledgements**

487 We thank our colleagues from the WSL, Dr. Ulf Büntgen, Dr. David Frank and Richard L. Peters,
488 for their help with ring-width and climate analyses, Dr. Alexander Bast for helping with
489 programming in R, and Anne Verstege for her continuous support in the dendro laboratory. Also we
490 wish to thank Vincenzo Crimi (Corpo Forestale, Bronte) for his logistical support during our
491 fieldwork on Mt. Etna, as well as the Corpo Forestale della Regione Siciliana (Distaccamento di
492 Bronte, Catania, Italy) for their permission to take samples.

493 We celebrate the memory of our colleague and friend, Sebastiano Cullotta, who tragically passed
494 away while this paper was in preparation. This work was supported by the Swiss National
495 Foundation (205321_143479).

496

497

498 **References**

499 Aiuppa A, Moretti R, Federico C, Giudice G, Gurrieri S, Liuzzo M, et al. Forecasting Etna
500 eruptions by real-time observation of volcanic gas composition. *Geology* 2007;35:1115-1118.

501 Akkemik Ü, D'Arrigo R, Cherubini P, Köse N, Jacoby G. Tree-ring reconstructions of precipitation
502 and streamflow for north-western Turkey. *Int J Climatol* 2008;28:173-183.

503 Allard P. Endogenous magma degassing and storage at Mount Etna. *Geophys Res Lett*
504 1997;24:2219-2222.

505 Allard V, Ourcival JM, Rambal S, Joffre R, Rocheteau A. Seasonal and annual variation of carbon
506 exchange in an evergreen Mediterranean forest in southern France. *Glob Chang Biol*
507 2008;14:714-725.

508 Aloui A. Recherches dendroclimatologiques en Kromirie (Tunisie). PhD. Thesis, Université d'Aix-
509 Marseille III, France. 1982.

510 Andronico D, Branca S, Calvari S, Burton M, Caltabiano T, Corsaro RA, et al. A multi-disciplinary
511 study of the 2002-03 Etna eruption: insights into a complex plumbing system. Bull Volcanol
512 2005;67:314-330.

513 Andronico D, Scollo S, Cristaldi A, Ferrari F. Monitoring the ash emission episodes at Mt. Etna: the
514 16 November 2006 case study. J Volcanol Geotherm Res 2009;180:123-134.

515 Aubert M. Practical evaluation of steady heat discharge from dormant active volcanoes: case study
516 of Vulcarolo fissure (Mount Etna, Italy). J Volcanol Geotherm Res 1999;92:413-429.

517 Battaglia M, Roberts C, Segall P. Magma intrusion beneath long valley caldera confirmed by
518 temporal changes in gravity. Science 1999;285:2119-2122.

519 Begum S, Nakaba S, Yamagishi Y, Oribe Y, Funada R. Regulation of cambial activity in relation to
520 environmental conditions: understanding the role of temperature in wood formation of trees.
521 Physiol Plant 2013;147:46-54.

522 Bolle HJ, Eckardt M, Koslowsky D, Maselli F, Melia Miralles J, Menenti M, et al. Mediterranean
523 land surface processes assessed from space. Regional Climate Studies; vol. XXVIII. Springer
524 Series; 2006.

525 Bonneville A, Gouze P. Thermal survey of Mount Etna volcano from space. Geophys Res Lett
526 1992;19:725-728.

527 Branca S, Coltelli M, De Beni E, Wijbrans J. Geological evolution of Mount Etna volcano (Italy)
528 from earliest products until the first central volcanism (between 500 and 100 ka ago) inferred
529 from geochronological and stratigraphic data. Int J Earth Sci 2008;97:135-152.

530 Briffa K, Schweingruber FH, Jones PD, Osborn TJ, Harris IC, Shiyatov SG, et al. Trees tell of past
531 climates: but are they speaking less clearly today? Philosophical Transactions of the Royal

532 Society of London B Biological Sciences 1996;353:65-73.

533 Briffa K, Melvin TM, Osborn TJ, Hantemirov RM, Kirilyanov AV, Mazepa VS, et al. Reassessing
534 the evidence for tree-growth and inferred temperature change during the Common Era in
535 Yamalia, northwest Siberia. *Quat Sci Rev* 2013;72:83-107.

536 Buntgen U, Frank DC, Nievergelt D, Esper. Summer temperature variations in the European Alps,
537 A.D. 755–2004. *J Clim* 2005;9:5606-5623.

538 Buntgen U, Frank DC, Schmidhalter M, Neuwirth B, Seifert M, Esper J. Growth/climate response
539 shift in a long subalpine spruce chronology. *Trees* 2006;20:99-110.

540 Buntgen U, Frank DC, Neuenschwander T, Esper. Fading temperature sensitivity of Alpine tree
541 growth at its Mediterranean margin and associated effects on large-scale climate reconstructions.
542 *Clim Change* 2012;114:651-666.

543 Burga C, Klötzli F. *Gebirge der Erde. Landschaft, Klima, Pflanzenwelt*. Stuttgart: Eugen Ulmer
544 GmbH & Co.; 2004.

545 Cai Q, Liu Y, Lei Y, Bao G, Sun. Reconstruction of the March-august PDSI since 1703 AD based
546 on tree rings of Chinese pine (*Pinus tabulaeformis* Carr.) in the Lingdong Mountain, southeast
547 Chinese loess Plateau. *Clim Past* 2014;10:509-521.

548 Camarero JJ, Manzanedo RD, Sánchez-Salguero R, Navarro-Cerrillo. Growth response to climate
549 and drought change along an aridity gradient in the southernmost *Pinus nigra* relict forests. *Ann*
550 *For Sci* 2013;70:769-780.

551 Campelo F, Nabais C, Freitas H, Gutiérrez E. Climatic significance of tree-ring width and intra-
552 annual density fluctuations in *Pinus pinaster* from a dry Mediterranean area in Portugal. *Ann For*
553 *Sci* 2006;64:229-238.

554 Carrer M, Nola P, Motta R, Urbinati C. Contrasting tree-ring growth to climate responses of *Abies*
555 *alba* toward the southern limit of its distribution area. *Oikos* 2010;119:1515-1525.

556 Certini G, Sanjurjo MJF, Corti G, Ugolini F. The contrasting effect of broom and pine on pedogenic
557 processes in volcanic soils (Mt. Etna, Italy). *Geoderma* 2001;102:239–254.

558 Cherubini P, Gartner BL, Tognetti R, Bräker OU, Schoch W, Innes J. Identification, measurement
559 and interpretation of tree rings in woody species from Mediterranean climates. *Biol Rev Camb*
560 *Philos Soc* 2003;78:119-148.

561 Christensen JH, Krishna Kumar K, Aldrian E, An SI, Cavalcanti IFA, de Castro M, et al. Climate
562 Phenomena and their Relevance for Future Regional Climate Change. In: *Climate Change 2013:*
563 *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of*
564 *the Intergovernmental Panel on Climate Change* (Stocker TF, Qin D, Plattner GK, Tignor M,
565 Allen SK, Boschung J, et al. (Eds.)). Cambridge University Press, Cambridge, United Kingdom
566 and New York, USA. 2013.

567 Cook ER. A time series approach to tree-ring standardization, PhD. Thesis, Laboratory of Tree-
568 Ring Research. Tucson: University of Arizona. 1985. Available:
569 <http://lrr.arizona.edu/sites/lrr.arizona.edu/files/bibliodocs/CookER-Dissertation.pdf>

570 Cook E, Kariukstis L. *Methods of dendrochronology - Applications in the environmental sciences.*
571 Dordrecht: Kluwer Academic Publishers; 1990.

572 Cook ER, Pederson N. Book- Chapter: Uncertainty, emergence and statistics in dendrochronology.
573 In: *Dendroclimatology, Developments in Paleoenvironmental Research* 2011;11:77-112.

574 Cotrufo MF, Alberti G, Inglema I, Marjanovic H, LeCain D, Zaldei A, et al. Decreased summer
575 drought affects plant productivity and soil carbon dynamics in a Mediterranean woodland.
576 *Biogeosciences* 2011;8:2729-2739.

577 Dazzi C. Environmental features and land use of Etna (Sicily - Italy). In: Arnalds O, Bartoli F,
578 Buurman P, Oskarsson H, Stoops G, Garcia-Rodeja E, editors. *Soils of Volcanic Regions in*
579 *Europe.* Heidelberg: Springer-Verlag Berlin; 2007.

580 Dobbertin M. Tree growth as indicator of tree vitality and of tree reaction to environmental stress: a
581 review. *Eur J For Res* 2005;124:319-333.

582 Doglioni C, Innocenti F, Mariotti G. On the geodynamic origin of Mt. Etna. *GNGTS - Atti del 17*
583 *Convegno Nazionale*, 2002.

584 Eckstein D, Aniol R. Dendroclimatological reconstruction of the summer temperatures for an alpine
585 region. *Mitt Forstl Bundes-Vers.anst Wien* 1981;142:391-298.

586 Egli, M, Mirabella, A, Nater M, Alioth L, Raimondi S. Clay minerals, oxyhydroxide formation,
587 element leaching and humus development in volcanic soils. *Geoderma* 2008;143:101-114.

588 Egli M, Mastrolonardo G, Seiler R, Raimondi S, Favilli F, Crimi V, et al. Charcoal and stable soil
589 organic matter as indicators of fire frequency, climate and past vegetation in volcanic soils of
590 Mt.Etna, Sicily. *Catena* 2012;88:14-26.

591 Fritts HC. *Tree Rings and Climate*. London: Academic Press Inc. Ltd; 1976.

592 Gamon JA, Field CB, Goulden ML, Friffin KL, Hartley AE, Joel G, et al. Relationships between
593 NDVI, canopy structure, and photosynthesis in three Californian vegetation types. *Ecological*
594 *Application* 1995;5:28-41.

595 García-Suárez AM, Butler CJ, Baille MG. Climate signal in tree-ring chronologies in a temperate
596 climate: A multi-species approach. *Dendrochronologia* 2009;27:183-198.

597 Gärtner H, Cherubini P, Fonti P, von Arx G, Schneider L, Nievergelt D, et al. A technical
598 perspective in modern tree-ring research - How to overcome dendroecological and wood
599 anatomical challenges. *J Vis Exp* 2015;97:e52337.

600 Gea-Izquierdo G, Cherubini P, Cañellas. Tree-rings reflect the impact of climate change on *Quercus*
601 *ilex* L. along a temperature gradient in Spain over the last 100 years. *For Ecol Manage*
602 2011;262(9):1807-1816.

603 Goetz SJ, Bunn AG, Fiske GJ, Houghton RA. Satellite-observed photosynthetic trends across
604 boreal North America associated with climate and fire disturbance. *Proc Natl Acad Sci USA*
605 2005;102:13521-13525.

606 Graumlich, LJ. Precipitation Variation in the Pacific Northwest (1675-1975) as Reconstructed from
607 Tree Rings. *Ann Assoc Am Geogr* 1987;77:19-29.

608 Griggs C, DeGaetano A, Kuniholm P, Newton M. A regional high-frequency reconstruction of
609 May-June precipitation in the north Aegean from oak tree rings, A.D. 1089-1989. *Int J Climatol*
610 2007;27:1075-1089.

611 Griggs C, Pearson C, Manning SW, Lorentzen B. A 250-year annual precipitation reconstruction
612 and drought assessment for Cyprus from *Pinus brutia* Ten. tree-rings. *Int J Climatol*
613 2014;34:2702-2714.

614 Grissino-Mayer HD. Evaluating crossdating accuracy: A manual and tutorial for the computer
615 program Cofecha. *Tree Ring Res* 2001;57(2):205-221.

616 Hermes K. Die Lage der oberen Waldgrenze in den Gebirgen der Erde und ihr Abstand zur
617 Schneegrenze (Kölner geographische Arbeiten 5). Universität Köln: Geographisches Institut;
618 1955.

619 Holmes RL. Computer-assisted quality control in tree ring dating and measurement. *Tree-ring Bull*
620 1983;43:69-78.

621 Hooper A, Prata F, and Sigmundson F. Remote sensing of volcanic hazards and their precursors.
622 *Proceedings of the IEEE* 2012;100(10):2908-2930.

623 Houlié N, Komorowski JC, de Michele M, Kasereka M, Ciraba H. Early detection of eruptive dykes
624 revealed by normalized difference vegetation index (NDVI) on Mt. Etna and Mt. Nyiragongo.
625 *Earth Planet Sci Lett* 2006;246:231-240.

626 Hughes MK. Aegean tree-ring signature years explained, *Tree-Ring Res* 2001;57:438-450.

627 Hughes MK. Dendrochronology in climatology - the state of the art. *Dendrochronologia* 2002;20(1-
628 2):95-116.

629 Jong SM. Imaging spectrometry for monitoring tree damage caused by volcanic activity in the Long
630 Valley caldera, California. *ITC Journal* 1998;1-10.

631 Köppen W. *Die Klimate der Erde. Grundriss der Klimakunde*. Berlin: Walter de Gruyter; 1923.

632 Körner C. A re-assessment of high elevation treeline positions and their explanation. *Oecologia*
633 1998;115:445-459.

634 Köse N, Akkemik Ü, Dalfes N. Tree-ring reconstructions of May-June precipitation for western
635 Anatolia. *Quat Res* 2011;75:438-450.

636 Köse N, Akkemik Ü, Güner HT, Dalfes HN, Frissino-Mayr HD, Özeren MS, et al. an improved
637 reconstruction of May-June precipitation using tree-ring data from western Turkey and its links
638 to volcanic eruptions. *Int J Biometeorol* 2013;57:691-701.

639 Kutner M, Nachtsheim C, Neter J, Li W. *Applied linear statistical models*. 5th ed. New York:
640 McGraw-Hill/Irwin; 2005.

641 Lebourgeois F, Levy G, Aussenac G, Clerc B, Willm F. Influence of soil drying on leaf water
642 potential, photosynthesis, stomatal conductance and growth in two black pine varieties. *Ann For*
643 *Sci* 1998;55:287-299.

644 Leonelli G, Pelfini M, Morra di Cella U. Detecting climatic treelines in the Italian Alps: The
645 influence of geomorphological factors and human impacts. *Phys Geogr* 2009;30(4):338-352.

646 Limm EB, Simonin KA, Bothman AG, Dawson TE. Foliar water uptake: a common water
647 acquisition strategy for plants of the redwood forest. *Oecologia* 2009;161:449-459.

648 Lüdeke MKB, Ramge PH, Kohlmaier GH. The use of satellite NDVI data for the validation of
649 global vegetation phenology models: application to the Frankfurt Biosphere Model. *Ecol Model*
650 1996;91:255-270.

651 Lulli L. Italian volcanic soils. In: Arnalds Ò, Bartoli F, Buurman P, Òskarsson H, Stoops G, Garcia-
652 Rodeja E, editors. Soils of volcanic regions in Europe. Berlin Heidelberg: Springer-Verlag;
653 2007. pp. 51–67.

654 Ma S, Baldocchi DD, Xu L, Hehn T. Inter-annual variability in carbon dioxide exchange of an
655 oak/grass savanna and open grassland in California. *Agric For Meteorol* 2007;147:157-171.

656 Maeda T, Takenaka H, Warkentin BP. Physical properties of allophane soils. *Adv Agron*
657 1977;29:229-264.

658 Maselli F. Monitoring forest conditions in a protected Mediterranean coastal area by the analysis of
659 multiyear NDVI data. *Remote Sens Environ* 2004;89:423-433.

660 Maselli F, Cherubini P, Chiesi M, Gilabert MA, Lombardi F, Moreno A, et al. Start of the dry
661 season as a main determinant of inter-annual Mediterranean forest production variations. *Agric*
662 *For Meteorol* 2014;194:197-206.

663 McNutt SR. Seismic monitoring and eruption forecasting of volcanoes: A review of the state-of-
664 the-art and case histories. Berlin Heidelberg: Springer-Verlag; 1996.

665 McNutt SR, Rymer H, Stix J. Synthesis of volcano monitoring. In: Encyclopedia of volcanoes.
666 Sigurdsson H, et al., editors. San Diego: Academic Press; 2000. pp. 1167-1185.

667 Mitchell TD, Carter TR, Jones PD, Hulme M, New M. A comprehensive set of high-resolution
668 grids of monthly climate for Europe and the globe: the observed record (1901-2000) and 16
669 scenarios (2001-2100). Working Paper 55. Norwich: Tyndall Centre for Climate Change
670 Research; 2004.

671 Navarro-Cerrillo RM, Sánchez-Salguero R, Manzanedo RD, Camarero JJ, Fernández Cancio A.
672 Site and age condition the growth responses to climate and drought of relict *Pinus nigra* subsp.
673 *salzmannii* populations in Southern Spain. *Tree Ring Res* 2014;70:145-155.

674 Olivar J, Bogino S, Spiecker H, Bravo F. Climate impact on growth dynamic and intra-annual
675 density fluctuations in Aleppo pine (*Pinus halepensis*) trees of different crown classes.
676 *Dendrochronologia* 2012;30:35-47.

677 O'Brian RM. A caution regarding rules of thumb for variance inflation factors. *Quality & Quantity*
678 2007;41:673-690.

679 Poli Marchese E, Grillo M. Primary succession on lava flows on Mt.Etna. In: Burga C.A., Klötzli
680 F., Grabherr G. (Eds.), *Gebirge der Erde. : Landschaft, Klima, Pflanzenwelt*. Stuttgart: Ulmer;
681 2004. pp. 291-300.

682 Reisser M. Soil analysis at Mount Etna to explain faster tree growth preceding a volcanic eruption.
683 Master Thesis. Department of Earth-System Sciences, University of Zurich. Switzerland; 2014.

684 Renault NL, Alvera B, Garcia-Ruiz JM. The snowmelt period in a Mediterranean high mountain
685 catchment: runoff and sediment transport. *Cuadernos de Investigacion Geografica* 2010;36:99-
686 108.

687 Scarascia-Mugnozza G, Oswald H, Piussi P, Radoglou K. Forests of the Mediterranean region: gaps
688 in knowledge and research needs. *For Ecol Manage* 2000;132:97-109.

689 Schweingruber FH. Flächenhafte dendroklimatische Temperaturrekonstruktionen für Europa.
690 *Naturwissenschaften* 1987;74:205-212.

691 Seim A, Büntgen U, Fonti P, Haska H, Herzig F, Tegel W, et al. Climate sensitivity of a
692 millennium-long pine chronology from Albania. *Clim Res* 2012;51:217-228.

693 Serre-Bachet F, Guiot J. Summer temperature changes from tree-rings in the mediterranean area
694 during the last 800 years. In: *Abrupt Climatic Change, Evidence and Implications*. NATO ASI
695 Series C, Mathematical and Physical Sciences, Vol. 216. The Netherlands: Reidel Publ. Co.;
696 1987. pp. 89-97.

697 Sherburn S, Scott B.J, Olsen J, Miller C. Monitoring seismic precursors to an eruption from the

698 Auckland volcanic field, New Zealand. New Zeal J Geol Geop 2007;50(1):1-11.

699 Shinohara, H. Excess degassing from volcanoes and its role on eruptive and intrusive activity. Rev
700 Geophys 2008;46:RG4005.

701 Sicali S, Barberi G, Cocina O, Musumeci C, Patanè D. Volcanic unrest leading to the July-August
702 2001 lateral eruption at Mt. Etna: Seismological constraints. J Volcanol Geotherm Res
703 2015;304:11-23.

704 Sparks RSJ, Biggs J, Neuberg JW. Monitoring Volcanoes. Science 2012;335:1310-1311.

705 Tardif J, Camarero JJ, Ribas M, Gutiérrez E. Spatiotemporal variability in tree growth in the central
706 Pyrenees: Climatic and site influences. Ecol Monogr 2003;73(2):241-257.

707 Tessier L. Dendroclimatologie et Ecologie de *Pinus sylvestris* L. et *Quercus pubescens* Willd. dans
708 le Sud-Est de la France. PhD. Thesis, Université d'Aix-Marseille III. 1984.

709 Todaro L, Andreu L, D'Alessandro CM, Gutiérrez E, Cherubini P, and Saracino A. Response of
710 *Pinus leucodermis* to climate and anthropogenic activity in the National Park of Pollino
711 (Basilicata, Southern Italy). Biol Cons 2007;137:507-519.

712 Tognetti R, Lombardi F, Lasserre B, Cherubini P, Marchetti M. Tree-ring stable isotopes reveal
713 twentieth-century increases in water-use efficiency of *Fagus sylvatica* and *Nothofagus* spp. in
714 Italian and Chilean mountains. PLoS One 2014;9:e113136.

715